

Improvement in protein content and leaf rust resistance in semi dwarf wheat varieties through breeding research

ABRAR YASIN B¹ AND RAM M²

Department of Genetics and Plant Breeding¹, Directorate of Research², Sam Higginbottom Institute of Agriculture, Technology and Sciences, Naini-Allahabad, U.P. (211007) India

Received: January, 2018; Revised accepted: April, 2018

ABSTRACT

India witnessed Green Revolution with the introduction of semi-dwarf wheat varieties viz., Kalyan Sona and Sonalika from CIMMYT Mexico in 1963. With an increase in the wheat yield per unit area, the country became self-sufficient in wheat production. On the other hand, leaf or brown rust, one of the dreaded diseases of wheat is more prevalent in India. Therefore, the protein content and leaf rust resistance were taken as the major parameters to determine the improvement in these traits through breeding research. The trial was laid out in randomized block design with four replications during rabi season 2013-14. The results indicated that K-816 contained the maximum protein content (12.40%), followed by K-9533 (12.30%) and PBW-343 (12.28%). However, their differences in mean values were non-significant over checks. Almost all latest wheat varieties (except HUU-318, Veeri, K-816, Kalyan Sona and Sonalika) have shown a good degree of resistance to leaf rust races. It may be concluded from the present investigation that there has been no significant improvement brought in protein content through breeding research. However, resistance to leaf rust has been appreciated.

Key words: Wheat breeding, protein content, leaf or brown rust resistance

INTRODUCTION

Green revolution made a remarkable progress in increasing wheat production in India. With the introduction of semi-dwarf wheat varieties, especially two selections viz., Kalyan Sona and Sonalika made out of CIMMYT breeding materials, production of wheat in India rose from 12.0 m.t. in 1965 to 76.3 m.t. in 1999-2000 (Nagarajan, 2005). This period is most famous an era of 'Green Revolution' in the Indian history. Both Kalyan Sona and Sonalika were ideal in plant type. They possess short plant stature, bold and amber grains, respond to higher doses of fertilizer application (120N: 60P: 50K) and irrigation management (5-6) without lodging. Due to the short maturity duration both fitted well in the "rice-wheat" cropping system which collectively brought "Wheat Revolution". In addition, these varieties were resistant to all the three wheat rust diseases viz; stem rust, leaf rust and stripe rust. As a result production of wheat tremendously increased per unit area and time (Choudary and Ali, 2008). After yield, grain quality, especially protein is a very important character and direly needed for human beings.

However, even after the green revolution, in India bulk of population below poverty line is a victim of protein malnutrition, because 70% of protein needed by humans is met from cereal grains. There has been no significant improvement made in the protein content through breeding research because all the newly released high yielding improved varieties are at par in protein content when compared with the first generation semidwarf wheat varieties i.e. Kalyan Sona and Sonalika. Among wheat rusts, brown rust is most widespread, black rust is restricted to Peninsular India, yellow rust to Northern and North-Western regions of India (Thind, 2005). Each has a number of physiological races such as 32 of stem rust; 24 of leaf rust and 20 of stripe rust. Cultivars that only have race-specific rust resistant genes lose their effective resistance and become susceptible within a few years of release (James, 2013). It is, therefore, desirable that germplasm exhibiting resistance through non-specific interaction should be used in breeding programs rather than germplasms exhibiting only specific interaction. The foundation varieties Kalyan Sona and Sonalika also broke down their

resistance within five years to the leaf (brown) and yellow (strip) rust. These were quickly replaced by new semi-dwarf wheat varieties. Hence, the present investigation was undertaken to generate data to derive concrete and conclusive evidence to prove the level of improvement in grain quality (protein) characteristics and the level of resistance incorporated to brown rust (*P. recondita*) prevalent in Eastern India by breeding researches over Kalyan Sona and Sonalika (Check).

MATERIALS AND METHODS

The material used for present investigation comprised of 15 wheat varieties including two check lines (Kalyan Sona and Sonalika) which were obtained from wheat genetic stock, Directorate of Research, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad. The field experiment (1x5m) was laid out in the randomized block design with four replications at Experimental Research Farm, Department of Genetics and Plant Breeding, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad. Each plot consisted of 4 rows, 1x5m plot size with plant geometry of 25x5cm. All the recommended cultural practices were followed during the crop growth period. For protein analysis, first primary grain samples were collected from each replication and from this primary sample, working grain sample lots were made which were subjected to grain protein estimation as per method suggested by Lowry *et al.* (1951). After observing the protein content percentage of each wheat variety, their protein estimation difference over the check was determined according to the following formula (Gomez and Gomez, 2010).

$$\bar{X}_1 - \bar{X}_2$$

Where \bar{X}_1 = Mean protein content of variety

\bar{X}_2 = Mean protein content of check

Protein percentage increase/decrease over the check was determined according to the following formula (Andrew, 2001).

$$\frac{\bar{X}_1 - \bar{X}_2}{\bar{X}_2} \times 100$$

Where \bar{X}_1 = Mean protein content of variety

\bar{X}_2 = Mean protein content of check

On the basis of CD (0.5%) values, the significant and non-significant performance of 13 semi-dwarf wheat varieties for grain protein content over two checks (Kalyan Sona and Sonalika) was determined by using the following formula (Gomez and Gomez, 2010):

$$\text{CD value} \pm \bar{X}_2 = \bar{X}_3$$

Where, \bar{X}_2 = Mean protein content of check

\bar{X}_3 = Estimated value

Estimated value was compared with mean protein value of variety, if estimated value recorded below the mean protein value of variety then mean performance of variety for protein content percentage is considered to be significant over check and if estimated value exceeds the mean value of variety then mean performance of variety for protein content percentage is considered to be non significant over check.

The infection of rust was checked and measured on each plot and on the basis of intensity of infection (percentage of leaf area infected) plants were grouped using the scale developed by U.S. Department of Agriculture (Herbert *et al.*, 1955) into highly resistant (HR-0% infection), resistant (Traces, 1-10%), moderate resistant (MR, 11-20%), moderate susceptible (MS, 21-30%), susceptibility (S, 31-50%), highly susceptible (HS, 51% to above).

RESULTS AND DISCUSSION

The data on mean estimation of protein percentage of 15 semi-dwarf wheat varieties including two checks (Kalyan Sona and Sonalika) have been summarized in Table 1. The perusal of the protein data indicated that K-816 contained the maximum amount of protein (12.40%) among the 15 varieties studied in this experiment including checks (Kalyan Sona (11.90%) and Sonalika (11.78%). However, the percentage increase in protein content of K-816 is hardly 4.20% over Kalyan Sona and 5.26% over Sonalika. The next high protein yielding variety was K-9533 (12.30%) followed by PBW-

343 (12.28%) but increase in the protein content of K-9533 and PBW-343 over Kalyan Sona is just 3.36% and 3.19% and 4.41% and 4.24% over Sonalika. Raj-3777 (-6.38%, -5.77%), UP-2594 (-5.88%, -4.92), HD-2009 (-4.78%, -3.82%), Raj-4037 and HD-2824 (-3.02%, -2.03%) were found inferior in respect of protein contents than Kalyan Sona and Sonalika (checks). It is evident (Table-1) that none of the high yielding semi-dwarf wheat varieties showed significant improvement in the protein content over Kalyan Sona and Sonalika. It appears that quality aspect has remained untouched by the wheat breeders. Harish (2001) reported that the protein content in Indian wheat

ranged between 11 and 12%, as against over 13% in Australian or North American wheat. Shashi *et.al.*, (2004) confirmed that wheat varieties from Punjab, Haryana, Uttar Pradesh, Bihar and Madya Pradesh have 11-12% protein content. Thus, our observations are in conformity with Harish (2001) and Shashi *et al.* (2004). Since 90% of semi-dwarf wheat varieties released in India are selection/introduction of elite lines evolved at CIMMYT, Mexico (Nagrajan, 2005). It appears that even at CIMMYT Mexico, no attempt was made to improve the protein content beyond 11 to 12% in the newly evolved wheat varieties.

Table 1: Comparative mean protein content of 15 semi dwarf wheat varieties including two checks

Name of Variety	Mean estimation of protein (%)	Protein estimation difference (%) over two checks		Percentage increase /decrease over check	
		K. Sona	Sonalika	K. Sona	Sonalika
PBW-343	12.28	+0.38	+0.50	3.19	4.24
PBW-373	11.66	-0.24	-0.12	-2.01	-1.00
HD-2733	11.81	-0.09	+0.03	-0.75	0.25
HD-2824	11.54	-0.36	-0.24	-3.02	-2.03
HD-2009	11.33	-0.57	-0.45	-4.78	-3.82
Veeri	11.63	-0.27	-0.15	-2.26	-1.27
HUW-510	11.73	-0.17	-0.05	-1.42	-0.42
HUW-318	11.62	-0.28	-0.16	-2.35	-1.35
K-816	12.40	+0.50	+0.62	4.20	5.26
K-9533	12.30	+0.40	+0.52	3.36	4.41
RAJ-3777	11.14	-0.76	-0.68	-6.38	-5.77
RAJ-4037	11.54	-0.36	-0.24	-3.02	-2.03
UP-2594	11.20	-0.70	-0.58	-5.88	-4.92
K.Sona (Check)	11.90				
Sonalika(Check)	11.78				
SE(x) ±	0.334				
CD (5%)	0.702				

*Significant over K.Sona, ** Significant over Sonalika and *** Significant over both the checks

Out of the three wheat rusts diseases, only leaf rust (*Puccinia recondita*) is a great problem in eastern U.P. Every year leaf rust infection starts by the first week of March and continues up to full March and cause about 10-15% loss in yield. Among the 15 varieties tested, only HUW-318 (21.24%), Veeri (21.25%) and K-816 (21.27%) showed moderate susceptibility to leaf rust (Table-2). Among two checks, Kalyan Sona also showed moderate susceptibility (21-23%) whereas Sonalika showed moderate resistance (11-16%) to leaf rust. Other wheat varieties were free from even traces of infection. It appears that all the newly semi-dwarf wheat varieties released after Kalyan Sona and Sonalika possess reasonably fair

degree of resistance to the leaf rust race flora prevalent in eastern U.P. This indicates that major attention has been paid to incorporate leaf rust resistance in the wheat breeding programme and during selection in breeding nursery as well as from entry to the promotion of new lines/genotypes from initial evaluation trial to uniform regional trials under AICWIP, the entries/lines having yield at par with check but resistant to the leaf rust were considered for promotion as a policy matter. A principal reason for the progress has been the number of trained individuals working on the problem in each wheat zone. This collective effort of plant pathologists and plant breeders in developing resistant cultivars and understanding disease

epidemiology has gradually reduced the magnitude and frequency of epidemics. Use of knowledge of pathogen variability generated thus far, formed the basis of a strategy for deployment of resistance genes in the field. The emergences of new races of rusts require continued efforts to deploy new resistance genes. It is obvious that gene management in the field has considerable promise for sustained control of rusts. The current status of rust resistance breeding involves both conventional

and molecular breeding approaches including QTL mapping. Efforts are being made since a long time to utilize wild relatives to develop novel germplasm. Wild species viz., *Ae. speltooides* (Dhakate, 2002), *Ae. triuncialis* (Aghaee *et al.* 2001), *Ae. ovata* (Dhaliwal *et al.* 2002), and *Ae. umbellulata* (Chunneja *et al.* 2008) have been exploited for leaf rust resistance. Useful genetic diversity is being continuously harnessed from wild relatives of wheat to diversify diseases resistance in wheat (Vinod *et al.*, 2009).

Table 2: Comparative data of leaf rust infection on 15 semi-dwarf wheat varieties including two checks

Name of Variety	Leaf Rust Infection in percentage	Status
PBW-343	F	Highly resistant (HR)
PBW-373	F	Highly resistant (HR)
HD-2733	F	Highly resistant (HR)
HD-2824	F	Highly resistant (HR)
HD-2009	F	Highly resistant (HR)
VEERI	21-25%	Moderate susceptibility(MS)
HUW-318	21-24%	Moderate susceptibility(MS)
HUW-310	F	Highly resistant (HR)
K-816	21-27%	Moderate susceptibility(MS)
K-9533	F	Highly resistant (HR)
Raj-4037	F	Highly resistant (HR)
Raj-3777	F	Highly resistant (HR)
UP-2594	F	Highly resistant (HR)
Kalyan Sona	21-23%	Moderate susceptibility(MS)
Sonalika	11-16%	Moderate resistant (MR)

F: Free from infection (Highly Resistance HR), 1-10% : (Traces) Resistance (R), 11-20% : Moderate Resistance (MR), 21-30% : Moderate Susceptibility MS), 31-50% : Susceptible (S), 51% to above : Highly Susceptible (HS)

On the basis of the present investigation, it may be concluded that no significant improvement has been achieved in protein percentage through breeding research. Protein content differences between the latest wheat varieties over the first series of semi-dwarf wheat

varieties Kalyan Sona and Sonalika has been very marginal. Of course, major attention has been paid to incorporate the resistance to leaf rust. Almost all the latest wheat varieties have shown a good degree of resistance to leaf rust disease.

REFERENCES

- Aghaee Sarbarzeh, M., Singh, Harjit and Dhaliwal, H.S. (2001) A microsatellite marker linked to leaf rust resistance transferred from *Aegilops triuncialis* into hexaploid wheat. *Plant Breeding* **120**: 259-261
- Andrew, J.Vicker (2001) The use of percentage change from baseline as an outcome in a controlled trial statistically inefficient: a simulation study. *BMC Medical Research Methodology*, <http://doi.org/10.1186/1471-2288-1-6>.
- Bains, N.S., Singh, S. and Dhillon, B.S. (2012) Enhanced utilization of plant genetic resources in crop improvement programme. *Indian Journal of Plant Genetic Resource* **25**: 52-62.
- Choudary, P.V.S. and Ali, S.M.A. (2008) Status paper on wheat. *Consortium of Indian Farmers Associations* pp: 3-6, http://www.indianfarmers.org/status_papers/6%20Status%20Paper%20on%20Wheat-Final.doc.

- Chunneja Kumar, P.S., Goel R.K., Aghaee Sabarzeh, M, Prashar, M and Dhaliwal, HS (2008) Transfer of leaf rust and stripe rust resistance from *Aegilops umbellulata* to bread wheat (*Triticum aestivum* L.). *Genetic Resource and Crop Evolution*. DOI 10.1007/s10722-007-9289-3.
- Dhakate, P.M. (2002) Genetic analysis of leaf rust and stem rust resistance(s) in interspecific derivatives of wheat (*Triticum aestivum* L.). *Ph.D. Thesis*, 93-pp, *Indian Agricultural Research Institute*, New Delhi.
- Dhaliwal, H.S., Chhuneja P., Goel R.K., Nayar, S.K., Kaur, R. and Singh, Harjit (2001) Introgression of novel sources of rust resistance from wild *Triticum* and *Aegilops* species into bread and durum wheat cultivars. *Indian Journal of Plant Genetic Resource* **14**: 153- 155.
- Harish, Damodarn (2001) Change in wheat export strategy mooted-cultivation of specific varieties conforming to global standards sought. *Business Line (International edition)*, *Financial Daily from The Hindu group of publications*.
- Herbert, K.H., Forrest, R.I. and David, C.S., (1955) *Methods of Plant Breeding*. *McGraw-Hill Publications*, USA pp: 143.
- James Kolmer (2013) Leaf rust of wheat-pathogen biology, variation and host resistance. *Forests* **4**: 70-84
- Koichi Fujita (2012) Green revolution in India and its significance in economic development: Implications for Sub-Saharan Africa. *JICA Research Institute* **15**: 5- 6.
- Gomez, K.A., and Gomez, A.A. (2010) *Statistical procedures for agriculture research*, Second edition. *John Walley & Sons, Inc UK*, pp:191
- Lowry, O.H., Rasebrough, N.J., Farr, A.L. and Randall, R.J. (1951) Protein measurement with the folin phenol reagent. *Journal of Biochemistry* **193**:165-175.
- Nagarajan, S. (2005) Can India produce enough wheat even by 2020? *Current Science* **89** (9):1467-1471.
- Shashi Madan, Rana, R.K. and Pawar, I.S. (2004) Evaluation of wheat varieties for quality characteristics. *National Journal of Plant Improvement* **6**:38-41
- Thind, T.S. (2005) *Diseases of field crops and their management*. *Daya Publishing House-Delhi*, pp: 1
- Vinod Raghu, B.R., Tomar, S.M.S., Sharma, J.B., Sinha Pallavi and Singh Bhanwar (2009) Exploitation of wild relatives for disease resistance in wheat. In Book of Abstracts. National Seminar on Designing crops for the changing climate, *Birsa Agricultural University, Ranchi*, pp. 227-22.