

Genetic variability and diversity in rice (*Oryza sativa* L.) landraces of Nagaland

M.K. MEENA¹, H.P. CHATURVEDI^{1*}, D. PURUSHOTAMA RAO¹ AND HARENDRRA VERMA²

¹Department of Genetics and Plant Breeding, School of Agricultural Sciences, Nagaland University, Medziphema- 797106

Received: September, 2025; Revised accepted: October, 2025

ABSTRACT

Rice is the principal food crop of Nagaland and plays a major role in tradition and culture of the ethnic tribes of the state. Leveraging the genetic variability present within the indigenous rice germplasm is necessary for further improvement of them. Hence, the present study was aimed to evaluate the genetic variability, genetic divergence and trait association of grain yield and its attributes in 22 rice landraces from the foothills of Nagaland. Observations were recorded for eight yield-contributing traits. Analysis of variance revealed significant variation for all the parameters under study. High genotypic and phenotypic coefficient variation was recorded for number of effective tillers and grain yield per hectare. While remaining yield traits showed medium to low range variation. In the present study, all yield and yield-related traits displayed high heritability. High heritability along with a high genetic advance as percentage of the mean was observed for plant height, number of effective tillers, panicle length, number of grains per plant and grain yield per hectare. Grain yield per hectare exhibited positive and significant correlations with plant height, number of effective tillers, and spikelet fertility, while days to 50 per cent flowering exhibited negative significant association with grain yield at both genotypic and phenotypic levels. All yield traits under study displayed significant positive correlations and positive direct effects to grain yield per hectare except days to 50 per cent flowering at both genotypic and phenotypic levels. The twenty-two genotypes were grouped into six clusters based on the relative magnitudes of D^2 values using Tocher's method. The present study suggested that the genotypes Boyoh, Ereima, Nyari, Rukhatang, Thevuru, Maso Tsuk, and Kemeny as the potential parents for hybridization programme to obtain desirable segregants.

Key words: Rice, Landraces, Variability, D^2 analysis, Correlation and Path analysis.

INTRODUCTION

Rice (*Oryza sativa* L.) is one of the world's most important staple cereal crops and is cultivated in 114 countries under diverse environmental conditions (Balakrishnan *et al.*, 2024). It forms an important food source for over 3.5 billion people contributing more than 20 per cent of total dietary calories. About 90 per cent of rice is produced and consumed in Asia. In India, rice is cultivated in a 43.79-million-hectare area with a production of 116.42 million tonnes in the year 2022-23 (Ratnam *et al.*, 2024). In Nagaland, a good number of local varieties and landraces with high genetic variability are maintained by the tribal farmers, which offer scope for further improvement through selection in spite of their low yield potential (Roy *et al.*, 2024). The improvement of these locally adapted genotypes for grain yield and quality attributing traits will reduce the pressure of increasing the land area under rice cultivation in Nagaland (Verma *et al.*, 2021). Nagaland has the richest pockets of rice germplasm in valleys as well as

in hilly areas. The landraces of rice exhibit large variations in morphology, thus signifying the vast genetic diversity among the cultivars (Lalrindiki *et al.*, 2024.). Assessment of genetic variation and diversity will create a benchmark for further improving the local landraces of rice in Nagaland. Grain yield is a complex trait contributed by various component traits and possesses low heritability. Therefore, selection for yield *per se* is ineffective due to its complex nature and, thus, indirect selection for yield attributing traits is more effective (Biswas *et al.*, 2024). Variability is the prerequisite for selection to be effective. The extent of phenotypic variability which is due to genotype governs the heritability. Heritability on the other hand indicates the extent to which trait variation is genetically determined (Krishnan *et al.*, 2024). For effective selection, traits not only possess high heritability but also high genetic advance.

Correlation analysis helps in understanding the interrelationship of different traits and thus provides basis of formulating selection scheme. Path coefficient analysis

Corresponding author's Email: hpchaturvedi68@gmail.com,
Medziphema, Nagaland

²ICAR Research Complex for NEH Region, Nagaland Centre,

further refines this by partitioning correlation coefficients into direct and indirect effects, providing deeper insight into trait relationships (Biswas *et al.*, 2024). Rice holds a significant place in the traditions of Nagaland's ethnic tribes and is deeply embedded in their cultural practices. The diverse rice landraces in Nagaland exhibit extensive phenotypic variation, adapting well to the wide-ranging hill environments. This diversity highlights the importance of identifying elite and novel variants that are well-suited to different agroclimatic conditions, offering valuable resources for future breeding efforts. In this study, 22 rice landraces grown in the foothills of Nagaland were characterized and subjected to variability and diversity studies. The information obtained from the study would be helpful for breeders for undertaking rice improvement programme in the foothills of Nagaland.

MATERIALS AND METHODS

Experiment location and Plant material

The study was conducted during the *Kharif* season-2023, at the Research Farm of the Department of Genetics and Plant Breeding, School of Agricultural Science, Nagaland University, Medziphema, India. A total of 22 rice landraces collected from various parts of Nagaland.

Experimental design and field layout

The study employed a randomized complete block design (RCBD) with three replications. Within each replication, there were twenty-two plots measuring 3m x 3m, spaced at intervals of 50 cm. Plant spacing within rows was 15 cm, while row spacing was 20 cm. Recommended agricultural practices were followed to raise the crop.

Data collection

A total of eight quantitative characters on 22 rice landraces genotypes were considered for experimentation. These characteristics were chosen based on descriptions and guidelines provided by PPV & FR in 2001 (DUS). For each

characteristic, data were gathered from five randomly sampled plants from each genotype and in each replication. The eight quantitative traits under the investigation were *viz.*, day to 50 per cent flowering (DF), days of maturity (DM), Plant height (PH), number of effective tillers per hill (TPH), Panicle length (PL), number of grains per plant (GP), Spikelet fertility (SF) and Grain yield per hectare (GYPH).

Statistical analysis

An analysis of variance (ANOVA) was conducted using the OPSTAT open-source software to evaluate the data. The phenotypic, genotypic, and environmental coefficients of variation were calculated following the methods outlined by Burton and De Vane (1953). Heritability estimations were derived according to Allard (1960). The potential genetic advance through selection was calculated based on the approach by Johnson *et al.*, (1955). Phenotypic and genotypic correlation coefficients were determined as suggested by Al-Jibouri *et al.*, (1958). The partitioning of the genotypic correlation coefficients into direct and indirect effects was done using the method proposed by Dewey and Lu (1959). Genetic diversity was estimated using D^2 statistic (Mahalanobis, 1936). The genetic distance ('D') between clusters was derived by taking the square root of the average D^2 values. Genotypes were grouped into clusters using Tocher's method based on the ascending order of their D^2 following Singh and Choudhury (1985).

RESULTS AND DISCUSSION

Analysis of variance

The results of the analysis of variance for the eight quantitative traits under investigation are presented in Table 1. The results indicated significant variation ($P < 0.05$) among the genotypes for all the traits suggesting presence of substantial amount of genetic variation among the evaluated genotypes. Comparable findings were also observed in studies conducted by Demeke *et al.*, (2023) and Asante *et al.*, (2019).

Table 1: Analysis of variance for eight characters in 22 genotypes of rice

Source of variance	Mean square		
	Replications	Genotypes	Error
Degree of freedom	2	21	42
Days to 50% flowering	97.059	193.900**	0.344
Days to maturity	59.066	358.801**	0.313
Plant height	80.01	1697.463**	0.239
Number of effective tillers	2.871	6.870**	0.046
Panicle length	34.911	23.472**	0.161
Number of grain per plant	44.492	1760.152**	0.122
Spikelet fertility	71.225	68.312**	0.289
Grain Yield per hectare	5.353	2.429**	0.043

Genetic Parameters of Variation

The success of crop breeding relies heavily on the genetic variability present in the population and the heritability of the traits. To

select desirable traits for trait modeling, a plant breeder must analyze the genetic variation within the existing population (Gupta *et al.*, 2020). The genetic variance components for yield and yield-related traits are presented in Table 2.

Table 2: Genetic parameters of yield and its related traits in 22 genotypes of rice

Character	Coefficient of variation		Heritability (Broad sense) %	Genetic advance as % of mean
	GCV %	PCV %		
Days to 50% flowering	8.953	8.961	99.81	18.427
Days to maturity	9.098	9.102	99.9	18.734
Plant height	16.284	16.285	99.8	33.543
Number of effective tillers	23.307	23.385	99.3	47.853
Panicle length	10.35	10.386	99.33	21.248
Number of grain per plant	17.32	17.321	99.99	35.678
Spikelet fertility	5.331	5.343	99.6	10.959
Grain yield per hectare	23.34	23.55	98.22	47.653

GCV= Genotypic coefficient of variation, PCV= Phenotypic coefficient of variation

The results of the present study revealed that the phenotypic coefficient of variance (PCV) was higher than the genotypic coefficient of variance (GCV) for all the traits. This might be due to influence of environmental effects. High genotypic and phenotypic coefficients of variation (greater than 20%) were observed for number of effective tillers and grain yield per hectare. This indicated that simple selection would be effective for these traits. Similar findings were also reported in other studies by Punyo and Chaturvedi (2020), Debsharma *et al.*, (2022) Sanku *et al.*, (2022) and Kangana Thakur *et al.* (2025). Mere presence of genetic variation may not indicate the effectiveness of selection unless the heritability and genetic advance are not known. In the present study, all yield and yield-related traits displayed high heritability (>60%), indicating a strong correlation between phenotypic and genotypic values. However, the success of selection depends not

only on heritability but also on genetic advance (Debsharma *et al.*, 2022). High heritability along with a high genetic advance as percentage of the mean was observed for plant height, number. of effective tillers, panicle length, number of grains per plant and grain yield per hectare. It could be inferred that these traits predominantly exhibited additive gene action, suggesting that simple selection methods could effectively improve them. High heritability along with medium genetic advance as percentage of the mean, was observed for days to flowering, maturity and spikelet fertility. These findings are supported by the observations made earlier by Srujana *et al.*, (2017), Thuy *et al.*, (2023), Deivanai *et al.*, (2025), H.M. Sachin Kumar *et al.*, (2025) and Islam *et al.*, (2025).

Correlation Studies

Understanding the strength of the

association between yield and its contributing traits is necessary for planning an effective selection program. The correlation coefficients (both genotypic and phenotypic) for grain yield and yield traits are presented in Table 3. Grain yield per hectare exhibited positive and significant correlations with plant height, number of effective tillers and spikelet fertility, while days to 50 per cent flowering exhibited negative significant correlation with grain yield at both

genotypic and phenotypic levels. This suggests that these traits are genetically linked or share common genetic factors contributing to yield. The strong correlations observed at both genotypic and phenotypic levels indicate that selecting these traits could effectively improve seed yield. Similar findings on the correlation of these traits with seed yield were also reported by Srujana *et al.*, (2017) and Thuy *et al.*, (2023).

Table 3: Genotypic and Phenotypic correlation of eight yield traits in 22 rice genotypes

Genotypic correlation	Characters	Days to 50% flowering	Days to maturity	Plant height	Number of effective tillers	Panicle length	Number of grain per plant	Spikelet fertility	Yield per hectare	Phenotypic correlation
	Days to 50% flowering	1	0.479*	-0.331	-0.591**	-0.352*	-0.2637	-0.209	-0.366*	
	Days to maturity	0.479*	1	-0.27	-0.358	-0.765**	-0.463*	-0.203	-0.123	
	Plant height	-0.331	-0.27	1	0.593**	0.201	0.144	0.388*	0.457*	
	No. of effective tillers	-0.591**	-0.358	0.593**	1	0.1967	-0.099	0.441*	0.503**	
	Panicle length	-0.352*	-0.765**	0.201	0.1967	1	0.295	-0.065	0.033	
	No. grain per panicle	-0.2637	-0.463*	0.144	-0.099	0.295	1	0.002	0.022	
	Spikelet fertility	-0.209	-0.203	0.388*	0.441*	-0.065	0.002	1	0.454**	
	Grain yield per hectare	-0.366*	-0.123	0.457*	0.503**	0.033	0.022	0.454**	1	

** = Significant at 1% level, * = Significant at 5% level

Path coefficient analysis

Path coefficient analysis was conducted at the genotypic level to differentiate the direct and indirect effects of various yield-contributing traits, as presented in Table 4. The traits, *viz.*, plant height, number of effective tillers and spikelet fertility under study displayed significant positive correlations and positive direct effects to grain yield per hectare. Therefore, emphasis should be placed on selecting these traits to

obtain improvement on grain yield. These findings are consistent with of the observations of Gupta *et al.*, (2020). The residual effect was estimated at 0.7873, indicating that the traits studied do not fully account for the variability observed, suggesting that additional relevant characters may exist outside those currently investigated. Similar results were reported by Girma *et al.*, (2018) and Sadimantara *et al.*, (2021).

Table 4: Direct and indirect effects of yield components on seed yield at genotypic level in 22 rice genotypes

Characters	Days to 50% flowering	Days to maturity	Plant height	No. of effective tillers	Panicle length	No. of grain per plant	Spikelet fertility	r_g with Yield per hectare
Days to 50% flowering	-0.1866	-0.0894	0.0619	0.1103	0.0658	0.0492	0.0391	-0.3664
Days to maturity	0.1040	0.2171	-0.0586	-0.0779	-0.1661	-0.1007	-0.0442	-0.1236
Plant height	-0.0645	-0.0526	0.1945	0.1155	0.0392	0.0281	0.0756	0.4571
No. of effective tillers	-0.1283	-0.0778	0.1288	0.2169	0.0427	-0.0217	0.0958	0.5037
Panicle length	-0.0221	-0.0480	0.0127	0.0123	0.0628	0.0185	-0.0041	0.0337
No. of grain per plant	-0.0076	-0.0133	0.0041	-0.0029	0.0085	0.0287	0.0000	0.0022
Spikelet fertility	-0.0614	-0.0595	0.1137	0.1293	-0.0191	0.0000	0.2926	0.4549

RESIDUAL EFFECT = 0.7873

Genetic Divergence Analysis

In the present study genetic divergence

among the 22 rice genotypes of Nagaland was studied using Mahalanobis' D^2 statistic. This method determines the statistical distance

between the genotypes and based on which a hierarchical classification of genotypes can be obtained by following different clustering methods. In the present investigation clustering was done by Tocher's method. These clusters assist plant breeders in identifying genetically diverse parents within the population (Rajappa *et al.*, 2024). The current study aims to pinpoint genetically diversified parents for hybridization programs. The twenty-two genotypes were

grouped into six clusters based on the relative magnitudes of D^2 values using Tocher's method. Among these clusters, Cluster I contained the highest number of genotypes (14), followed by Cluster III, which included four genotypes. Clusters II, IV, V and VI each contained a single genotype. The distribution of genotypes across the four clusters is presented in Table 5, representing a high degree of genetic divergence among the genotypes.

Table 5: Clustering pattern of 22 rice genotypes on the basis of Tocher's method

Cluster	No. of Genotypes in Cluster	List of Genotypes
Cluster1	14	Aongsho, Malanken, Bhalum - I, Manabe, Manen Tsuk, Chali Yoh, Doulong, Moro Etyo, Hyung Yoh, Samro Yoh, Semmeki, Koyapvu Tsuk, Shangya, Lamjet
Cluster2	1	Engcha Yoh
Cluster3	4	Kemenya, Rukhatang, Maso Tsuk, Thevuru
Cluster4	1	Nyari
Cluster5	1	Ereima
Cluster6	1	Boyoh

In D^2 analysis, cluster distance indicates the diversity between clusters. The inter-cluster distances measure the relatedness and genetic relationships among genotypes across different groups. Average intra and inter-cluster distances among the six clusters in rice landraces is presented in Table 6. In this study, cluster IV and VI exhibited the largest average inter-cluster distance of 366.49 followed by cluster II and IV (295.69), cluster III and IV (281.52) and Cluster V and VI (211.67). A greater inter-cluster distance suggests that the genotypes within these clusters are more diverse. Conversely, a smaller inter-cluster distance indicates a closer relationship among the genotypes in those clusters, typically reflecting similar gene complexes (Padmakumar *et al.*, 2024). The maximum inter-cluster distance points to significant diversity among genotypes across the groups (Singh *et al.*, 2024). The promising

genotypes identified in these clusters could serve as potential parents for breeding. Crosses between genotypes from distinct clusters are believed to have the highest potential for producing beneficial recombinants, which can enhance yield component traits (Singh *et al.*, 2024). Based on the clustering pattern in the present study suggests the genotypes Boyoh, Ereima, Nyari, Rukhatang, Thevuru, Maso Tsuk, and Kemeny were identified as potential parents and suitable candidates for hybridization programme. Geographic distribution significantly influences genetic divergence and relationships among the genotypes across different environments (Rao and Chaturvedi, 2024). Geographic factors can lead to the development of unique genotypes with specific adaptations, resulting in diverse clusters and varying distances between them (Rao and Chaturvedi, 2024).

Table 6: Average intra and inter- cluster incidents among four clusters in 22 rice genotypes

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6
Cluster 1	85.07	118.15	140.41	231.57	167.45	213.5
Cluster 2		0	200.81	295.69	170.14	108.54
Cluster 3			83.84	126.54	127.2	281.52
Cluster 4				0	180.61	366.49
Cluster 5					0	211.67
Cluster 6						0

Cluster mean values are presented in Table 7. Among the clusters, Cluster VI exhibited the highest mean value for the number of grains per plant (198.10), with a lower value of yield per hectare (2.83 tons). Cluster IV, with the highest

yield per hectare (4.23 tons), was also promising, despite showing moderate values for other traits. Clusters I and IV are ideal due to their respective high mean plant heights (158.06 cm and 143.73 cm).

Table 7: Cluster wise mean value of eight characters in 22 rice genotypes

	Days to 50% flowering	Days to maturity	Plant height	No. of effective tillers	Panicle length	No. of grain per plant	Spikelet fertility	Yield per hectare
Cluster 1	87.85	117.13	158.06	6.75	27.58	145.35	89.49	4.00
Cluster 2	81.87	109.27	139.07	7.40	30.37	171.60	89.10	3.30
Cluster 3	97.52	125.87	135.33	6.60	24.85	113.56	91.98	3.48
Cluster 4	82.33	143.73	114.73	6.10	22.83	87.20	86.83	4.23
Cluster 5	101.10	126.67	95.63	3.93	29.90	130.57	81.93	3.80
Cluster 6	88.53	120.30	109.77	4.10	23.87	198.10	86.30	2.83

Thus, Clusters IV and VI appeared to be significant for selection of parents for inclusion in hybridization programme to obtain desirable segregants. The character plant height (32.03%) and number of grains per panicle (60.17%) contributed maximum towards the divergence (Table 8).

Table 8: Contribution of each character of 22 rice genotypes towards the divergence

Source	Contribution %
Days to 50% flowering	0.87%
Days to maturity	3.90%
Plant height	32.03%
Number of effective tillers	0.43%
Panicle length	0.00%
Number of grain per panicle	60.17%
Spikelet fertility	2.60%
Yield per hectare	0.00%

CONCLUSION

In the present study, traits viz., plant

height, number of effective tillers, panicle length, number of grains per plant and grain yield per ha exhibited high heritability coupled with high genetic advance suggesting selection based on these traits would be effective for improving grain yield. Plant height, number of effective tillers and spikelet fertility under study displayed significant positive correlations and positive direct effects to grain yield per hectare. Therefore, emphasis should be placed on selecting these traits to obtain improvement on grain yield. The genotypes viz., Boyoh, Nyari, Ereima, Kemenya, Rukhatang, Maso Tsuk, and Thevuru exhibited the highest number of productive tillers per plant, longer panicles, a higher percentage of spikelet fertility, and high yield. However, these genotypes also showed high inter-cluster distance indicating their genetic divergence. Therefore, they are considered divergent genotypes and are recommended for use in heterosis breeding programs.

REFERENCES

Al-Jibouri, H., Miller, P.A. and Robinson, H.F. (1958) Genotypic and environmental variances and covariances in an upland cotton cross of interspecific origin. *Agronomy Journal* **50**(10): 633–636.

Allard, R.W. (1961) *Principles of Plant Breeding*. John Wiley & Sons, New York.

Asante, M.D., Adjah, K.L. and Annan-Afful, E. (2019) Assessment of genetic diversity for grain yield and yield component traits in some genotypes of rice (*Oryza sativa* L.). *Journal of Crop Science and Biotechnology* **22**(2): 123–130.

Balakrishnan, C., Kumar, A., Raj, R., Verma, V.K., Touthang, L., Kumar, R. and Mishra, V.K. (2024) Exploring genetic diversity, population structure and stability for yield-related traits in rice germplasm of northeastern India. *Genetic Resources and Crop Evolution* **1**–21.

Biswas, A., Adhikari, A., Adhikari, S., Paul, A. and Ghosh, P. (2024) Assessment of variability and interrelationship between yield and yield-related traits towards divergence in rice (*Oryza sativa* L.) landraces. *The Nucleus* **1**: 1–16.

Burton, G.W. and De Vane, D.E. (1953) Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. *Agronomy Journal* **45**: 478–481.

Debsharma, S.K., Syed, M.A. and Ali, M.H. (2022) Harnessing on genetic variability and diversity of rice (*Oryza sativa* L.) genotypes based on quantitative and qualitative traits for desirable crossing materials. *Genes* **14**(1): 10.

Deivanai, A., Dhanapriya, M., Dhivyalakshmi, M. and Chitra, S. (2025) Studies on genetic variability in rice (*Oryza sativa* L.) genotypes under aerobic condition. *International Journal of Botany Studies* **10**(1): 56–58.

Demeke, B., Dejene, T. and Abebe, D. (2023) Genetic variability, heritability, and genetic advance of morphological, yield-related, and quality traits in upland rice (*Oryza sativa* L.) genotypes at Pawe, northwestern Ethiopia. *Cogent Food & Agriculture* **9**(1): 2157099.

Dewey, D.R. and Lu, K. (1959) A correlation and path-coefficient analysis of components of crested wheatgrass seed production. *Agronomy Journal* **51**(9): 515–518.

Girma, B.T., Kital, M.A., Banje, D.G., Biru, H.M. and Serbessa, T.B. (2018) Genetic variability study of yield and yield-related traits in rice (*Oryza sativa* L.) genotypes. *Advances in Crop Science and Technology* **6**(4): 381.

Gupta, S., Upadhyay, S. and Koli, G.K. (2020) Trait association and path analysis studies of yield attributing traits in rice (*Oryza sativa* L.) germplasm. *International Journal of Bio-resource and Stress Management* **11**(6): 508–517.

H.M., Sachin Kumar, Dushyantha Kumar, B.M., Krishnamurthy, S.L., Pattanashetti, S.K., Girijesh, G.K. and Halingali, B.I. (2025) Exploring the dynamics of genetic parameters: genetic variability, heritability and character association in rice (*Oryza sativa* L.). *Journal of Advances in Biology & Biotechnology* **28**(2): 52–58.

Islam, M.Z., Chakrabarty, T., Akter, Khalequzzaman, M., Prince, M. F. R. K., Pittendrigh, B. R and Ali, M. P. (2025) Genetic variability, correlation and path coefficient analysis of phenotypic traits and genetic diversity of Aman rice landraces (*Oryza sativa* L.). *Scientific Reports* **15**: 18606.

Johnson, H.W., Robinson, H.F. and Comstock, R.E. (1955) Estimates of genetic and environmental variability in soybeans. *Agronomy Journal* **47**(7): 314–318.

Kangkana Thakur, Sofia Banu and Manoj Kumar Sarma (2025) Genetic variability and diversity using SSR marker in indigenous winter rice germplasm of Assam. *Annals of Plant and Soil Research* **27**(1): 32-40

Krishnan, V., Sivarajani, V., Tamilzharasi, M. and Anandhan, T. (2023) Characterization of morpho-phenological traits in the traditional landraces of rice. *Electronic Journal of Plant Breeding* **14**(1): 234–245.

Lalrindiki, H., Kigwe Seyie, Chaturvedi, H.P. and Harendra Verma 2024. Genetic diversity analysis in some upland rice (*Oryza sativa* L.) landraces of Nagaland using grain quality and yield attributing traits. *Annals of Plant and Soil Research* **26**(4): 725-730

Mahalanobis P.C. (1936) On the generalized distance in statistics. *Proc. Nat. Inst. Sci. India (B)* **2**: 49-55.

Padmakumar, V., Ravi, D., Rupam, B., Kanwar, S., Ramakrishna, R., Prasad, K.V.S.V. and Jones, C.S. (2024) Genetic variation for grain yield and straw quality traits in 55 diverse rice varieties in Assam. *Animal Nutrition and Feed Technology* **24**(1): 93–103.

Rajappa, P.V., Anisha, A., Krishnaraj, P.U., Hanumaratti, N.G., Bagewadi, B., Nawalagatti, C.M. and Madhusudhana, R. (2024) Assessment of genetic diversity in rabi sorghum landraces using D² statistics. *Journal of Cereal Research* **16**(1): 92–101.

Rao, D.P. and Chaturvedi, H.P. (2024) Studies on genetic divergence in foxtail millet (*Setaria italica* (L.) P. Beauv.) genotypes grown over multi season at the Nagaland ecosystem. *Current Agriculture Research Journal* **12**(2): 941–957.

Ratnam, T.V., Kumar, B.R., Rao, L.V., Srinivas, T. and Kumar, A.A. (2024) Assessment of genetic variability, character association and path analysis for yield

and quality traits in zinc and iron-rich landraces of rice. *Agricultural Science Digest* **44**(2): 289–294.

Rinya Punyo and H. P. Chaturvedi (2020) Genetic variability, correlation and path coefficient studies in rice (*Oryza sativa* L.) genotypes. *Annals of Plant and Soil Research* **22**(3): 260-263.

Roy, S., Patra, B.C., Kumar, J., Sar, P., Jogi, U.S., Konyak, Z. and Bansal, K.C. (2024) Ethnolinguistic associations and genetic diversity of rice landraces in Nagaland, India. *Plants, People, Planet* **6**(2): 452–469.

Sadimantara, G.R., Yusuf, D.N., Febrianti, E., Leomo, S. and Muhidin, M. (2021) The performance of agronomic traits, genetic variability, and correlation studies for yield and its components in some red rice (*Oryza sativa*) promising lines. *Biodiversitas Journal of Biological Diversity* **22**(9): 3994–4001.

Sanku, G., Hepziba, S.J., Sheeba, A., Hemalatha, G. and Senthil, K. (2022) Genetic variability and relatedness among yield characters in rice landraces and improved varieties. *Electronic Journal of Plant Breeding* **13**(3): 932–939.

Singh Yadav, S.P., Bhandari, S., Ghimire, N.P., Mehata, D.K., Majhi, S.K., Bhattarai, S. and Bhujel, S. (2024) Genetic variability, character association, path coefficient, and diversity analysis of rice (*Oryza sativa* L.) genotypes based on agromorphological traits. *International Journal of Agronomy* **1**: 9946332.

Singh, R.K. and Chaudhary, B.D. (1985) Biometrical methods in quantitative genetic analysis. *Haryana Journal of Horticultural Sciences* **12**(2): 151–156.

Srujana, G., Suresh, B.G., Lavanya, G.R., Ram, B.J. and Sumanth, V. (2017) Studies on genetic variability, heritability, and genetic advance for yield and quality components in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry* **6**(4): 564–566.

Thuy, N.P., Nhu, T.T.T. and Trai, N.N. (2023) Genetic variability for micronutrients under an aerobic condition in local landraces of rice from different agro-ecological regions of Karnataka, India. *Biodiversitas Journal of Biological Diversity* **24**(1): 1–10.

Verma, H., Sharma, P.R., Chücha, D., Walling, N., Rajesha, G., Sarma, R.N. and Kandpal, B.K. (2021) Genetic characterization of local adaptable rice landraces of Nagaland, India. *Indian Journal of Plant Genetic Resources* **34**(2): 173–184.