

EFFECT OF PHOSPHORUS AND MOLYBDENUM ON YIELD AND NUTRIENT UPTAKE OF FABA BEAN IN ALLUVIAL SOIL

MUNNA LAL¹, ANIL KUMAR PAL², MAHESH CHAND AGRAWAL, K. USHA RANI³. D. SUMA CHANDRIKA¹ AND ABHAY PRATAP SINGH

Department of Agricultural Chemistry and Soil Science, Raja Balwant Singh College Bichpuri, Agra (U.P.)-283105

Received: November, 2015; Revised accepted March, 2016

ABSTRACT

An experiment was conducted at Bichpuri (Agra) during rabi seasons of 2008-09 and 2009-10 to study the impact of phosphorus and molybdenum on yield and nutrient uptake by faba bean (*Vicia faba*) in alluvial soil. The experiment was laid out in randomized block design with four levels of phosphorus (0, 30, 60 and 90 kg P₂O₅ ha⁻¹) and four levels of molybdenum (0, 1, 2, and 3 kg ha⁻¹). The results revealed that the faba bean responded significantly upto 90 kg P₂O₅ ha⁻¹ and increased the grain and straw yield by 13.4 and 15.7 percent, respectively, over control. Application of 3 kg Mo ha⁻¹ proved superior to the control with respect to grain and straw yields and increased the grain and straw yields by 9.0 and 8.4 percent, respectively. The phosphorus levels significantly increased the nitrogen, phosphorus, potassium and molybdenum uptake by faba bean over control. Molybdenum application also influenced the utilization of nitrogen, P, K and Mo and the more beneficial effect was observed with highest level of molybdenum (3 kg ha⁻¹). The status of nutrients in post harvest soil improved significantly with P and Mo levels over the control.

Key words: Phosphorus, molybdenum, yield, nutrient uptake, faba bean.

INTRODUCTION

Vicia faba, also known as the broad bean, fava bean or faba bean or field bean is a species of bean (Fabaceae). In addition to its use as food for human and livestock, faba bean plays a critical role in some agricultural systems due to the ability of the nitrogen-fixing bacteria, it harbors to fix the atmospheric nitrogen under a broad spectrum of environmental conditions. This facilitates diversification of the agro ecosystem in both time (via crop rotations) and in space (via intercrops), which also may indirectly enhance associated diversity of wild flora and fauna, as well as soil microbes, which may in turn impact the sustainability of agricultural systems. Molybdenum is required for growth of most biological organisms including plants and animals which play an important role in the process of *Rhizobium* symbiosis. Molybdenum is a constituent of the nitrogenase enzyme and every bacterium which fixes nitrogen needs molybdenum during the fixation process. Molybdenum has a positive effect on yield quantity, quality and nodule forming in legume crops (Singh *et al.* 2014). Phosphorus is

one of the most important elements significantly affecting plant growth and metabolism. It is, along with N, a major yield limiting nutrient. Phosphorus may be a critical constraint of legumes under low nutrient environments because there is a substantial need for P in the N₂ fixation process. However, in legumes, the high requirement for P is consistent with the involvement of P in high rates of energy transfer that take place in the nodule. Under P shortage conditions, legumes may lose the distinct advantage of an unlimited source of symbiotic N. In addition, phosphorus has also an enhancing impact on plant growth and biological yield through its importance as energy storage and transfer necessary for metabolic processes (Srivastava *et al.*, 1998 and Singh *et al.*, 2014). Phosphorus addition increased the efficiency of plants to photosynthesis, enhances the activity of rhizobia and increases the number of branches and pod/plants, consequently greater yield of pea, lupine, faba bean and groundnut. The present study was undertaken to evaluate the effect of phosphorus and molybdenum on faba bean crop in alluvial soil.

¹CRIDA, Hyderabad, ²Project Officer, Chhandiwara

MATERIALS AND METHODS

The field experiments were conducted at the Agriculture Research farm of R.B.S. College Bichpuri, Agra (located in semi arid or gray steppe arid region of South-Western Uttar Pradesh. the intersect of 27.2° N attitude and 77.9° E longitude), during two consecutive rabi seasons of 2008-09 and 2009-10 on sandy loam soil. The soil had EC 0.16 dSm⁻¹, pH 8.4, organic carbon 3.4 g kg⁻¹, available N 170, P 9.4, K 111 kg ha⁻¹, and molybdenum 0.05 mg kg⁻¹. The experiment was laid out in randomised block design with four levels of phosphorus (control, 30, 60 and 90 kg P₂O₅ ha⁻¹) and four levels of molybdenum (control, 1.0, 2.0 and 3 kg ha⁻¹) with three replications. The recommended doses of N and K @ 25 and 60 kg K₂O ha⁻¹, respectively were applied as urea and muriate of potash. Phosphorus and molybdenum were supplied through single super-phosphate and ammonium molybdate as per treatments. The faba bean was sown on October 25, 2008 and October 27, 2009 and irrigated at the proper time as judged by the appearance of soil and crop. The weeds were eradicated time to time from the crop. The crop was harvested on maturity. The grain and straw samples were analysed for N content by Kjeldahl method (Jackson 1973). Grain and straw samples were digested in di acid (HNO₃, HClO₄) and the digest were analysed for phosphorus by vanado molybophosphoric acid yellow colour method, K by flame photometer and Mo by atomic absorption spectrophotometer (Jackson 1973). The uptake of nutrients was calculated using the yield data in conjunction

with their respective contents. The soil samples collected after harvest were analysed for organic carbon, available N, P, K and Mo as per procedures suggested by Jackson (1973).

RESULTS AND DISCUSSION

Yield

The grain and straw yields of faba bean increased significantly by levels of phosphorus over control (Table 1). Application of 90 kg P₂O₅ ha⁻¹ recorded significantly higher yields of grain and straw over control. The increases in grain and straw yields with 90 kg P₂O₅ ha⁻¹ were 13.4 and 15.7%, respectively over control. The increase in yield may be attributed to the effective metabolic activities coupled with increased rate of photosynthesis leading to better translocation of nutrients to sink. Similar results were reported by Nusakho Nyekha *et al.* (2015) in green gram and Singh *et al.* (2016) in lentil. Pooled data showed 9.0 and 84% increase in grain and straw yield of faba bean (Table 1) with the application of 3 kg Mo ha⁻¹ over the control which might be owing to better nutritional environment in term of increased nitrogen fixation and increased plant growth. Molybdenum is known to be essential for N₂-fixation by rhizobia in legumes, being a component of nitrate reductase enzyme which controls the reduction of inorganic NO₃ and helps in fixing nitrogen as NH₃, as a result occurs increased nodulation, growth, grain and straw yield. These results confirm the findings of Kumawat *et al.* (2009) and Singh *et al.* (2014).

Table 1: Effect of Phosphorus and molybdenum on yield and uptake of N, P, K (kg ha⁻¹) and Mo (g ha⁻¹) by faba bean (mean of two years)

| Treatment | Yield (q ha ⁻¹) | | Uptake of nutrients | | | |
|-----------------------------------|-----------------------------|-------|---------------------|------|-------|-------|
| | Grain | Straw | N | P | K | Mo |
| Phosphorus (kg ha ⁻¹) | | | | | | |
| 0 | 35.76 | 30.33 | 139.1 | 27.0 | 87.9 | 201.9 |
| 30 | 37.24 | 31.93 | 150.4 | 30.5 | 94.6 | 231.8 |
| 60 | 39.53 | 33.83 | 164.1 | 34.2 | 103.5 | 259.0 |
| 90 | 40.57 | 35.10 | 172.5 | 38.2 | 108.7 | 294.7 |
| SEm± | 0.16 | 0.12 | 2.25 | 0.45 | 0.84 | 6.46 |
| CD (P = 0.05) | 0.44 | 0.35 | 6.54 | 1.26 | 2.43 | 18.10 |
| Molybdenum(kg ha ⁻¹) | | | | | | |
| 0 | 36.71 | 31.43 | 145.0 | 27.3 | 91.7 | 177.1 |
| 1 | 37.65 | 32.31 | 152.7 | 30.4 | 96.5 | 218.5 |
| 2 | 38.73 | 33.07 | 160.1 | 34.8 | 100.4 | 270.6 |
| 3 | 40.01 | 34.38 | 168.3 | 37.3 | 106.1 | 321.3 |
| SEm± | 0.16 | 0.12 | 2.25 | 0.45 | 0.84 | 6.46 |
| CD (P = 0.05) | 0.44 | 0.35 | 6.54 | 1.26 | 2.43 | 18.10 |

Nutrients Uptake

The uptake of nitrogen by faba bean crop increased significantly with increase in the levels of phosphorus compared to control. Application of 90 kg P₂O₅ ha⁻¹ resulted in significantly higher nitrogen uptake by the crop which may be attributed to increased N content coupled with higher yield (Singh *et al.* 2016). The uptake of nitrogen was significantly higher with increase in the levels of molybdenum over control and maximum value was recorded with 3 kg Mo ha⁻¹. This increase in N uptake may be attributed to the role of molybdenum in the process of *Rhizobium* symbiosis. Molybdenum, which is a constituent of nitrogenase enzyme, fixes nitrogen during the fixation process. Kushwaha (1999) and Singh *et al.* (2014) reported similar results. Increase in phosphorus levels significantly increased the phosphorus uptake by the crop over control and lower levels of phosphorus. The maximum enhancement in phosphorus uptake was recorded with 90 Kg P₂O₅ ha⁻¹. Similar findings were also recorded by Singh *et al.* (2016). The utilization of phosphorus by the crop significantly increased with increasing levels of molybdenum compared to control and the maximum value was recorded at 3 Kg ha⁻¹. Similar observations were also recorded by Kumawat *et al.* (2009). The uptake of potassium by the crop increased significantly with 60 and 90 kg ha⁻¹ compared to control. The

uptake of potassium did not show any significant change with 30 kg P₂O₅ ha⁻¹. Comparatively higher potassium utilization was recorded with 90 kg P₂O₅ ha⁻¹. The increase in potassium uptake might be due to increased potassium content and yield of faba bean with phosphorus levels. Our findings are in agreement with those of Singh *et al.* (2014). The uptake of potassium increased significantly with increase in the levels of molybdenum in comparison to control. The more beneficial effect on potassium uptake was noted with highest level of molybdenum (3 kg Mo ha⁻¹). Higher values of potassium uptake with molybdenum application are apparently the result of favourable effect on grain and straw production. Similar results were also noted by Srivastava and Ahlawat (1995) and Alben Awomi *et al.* (2012). The utilization of molybdenum increased with increase in levels of phosphorus over control and the maximum values were recorded with 90 kg P₂O₅ ha⁻¹. This increase is attributed to enhanced faba bean production and an increase in molybdenum content. A further study (Table 1), reveals that the molybdenum uptake by the crop increased significantly with increasing levels of molybdenum as compared to control. The maximum enhancement in molybdenum uptake was recorded with highest level of molybdenum (3 kg ha⁻¹). Similar findings were reported by Kushwaha, (1999).

Table 2: Effect of P and Mo levels on fertility status of post harvest soil (mean of 2 years)

| Treatment | Org. C (g kg ⁻¹) | Avail. N (kg ha ⁻¹) | Avail. P (kg ha ⁻¹) | Avail. K (kg ha ⁻¹) | Avail. Mo (mg kg ⁻¹) |
|-----------------------------------|------------------------------|---------------------------------|---------------------------------|---------------------------------|----------------------------------|
| Phosphorus (kg ha ⁻¹) | | | | | |
| 0 | 3.1 | 162.0 | 8.8 | 101.0 | 0.04 |
| 30 | 3.4 | 163.8 | 9.1 | 103.0 | 0.04 |
| 60 | 3.5 | 170.1 | 10.1 | 105.0 | 0.05 |
| 90 | 3.6 | 173.5 | 11.6 | 107.0 | 0.05 |
| SEm± | 0.011 | 0.63 | 0.32 | 0.93 | 0.006 |
| CD (P = 0.05) | 0.030 | 1.76 | 0.88 | 2.60 | NS |
| Molybdenum(kg ha ⁻¹) | | | | | |
| 0 | 3.3 | 163.3 | 9.1 | 102.5 | 0.03 |
| 1 | 3.4 | 164.5 | 9.4 | 103.5 | 0.04 |
| 2 | 3.4 | 169.0 | 10.0 | 104.4 | 0.05 |
| 3 | 3.5 | 171.0 | 10.3 | 105.1 | 0.06 |
| SEm± | 0.011 | 0.63 | 0.32 | 0.93 | 0.006 |
| CD (P = 0.05) | 0.030 | 1.76 | 0.88 | NS | 0.017 |

Soil fertility

Organic carbon content in post harvest soil increased from 3.1 g kg⁻¹ at control to 3.6 g kg⁻¹ with 90 kg P₂O₅ ha⁻¹. Similarly, Mo

application also enhanced the organic carbon content in soil. The minimum value of available N content in soil was recorded in control, which may be ascribed to greater utilization of N by

faba bean. Molybdenum application also improved the status of available N in soil as compared to control (Singh *et al.* 2014). There was a significant build-up of available P in soil with P application and maximum value was recorded with 90 kg P₂O₅ ha⁻¹ (Singh *et al.* 2016). The amount of available P in soil was lowest in control, which increased with Mo application. Similar results were reported by Singh *et al.* (2014). At harvest, the amount of available K was depleted from the initial value of 110 kg ha⁻¹ to 101 kg ha⁻¹. Available K content increased with various levels of Mo and maximum value was recorded with 3 kg Mo ha⁻¹.

Available Mo was the lowest in the control (Table 2). The concentration of available Mo increased significantly with 3 kg Mo ha⁻¹. This increase may be ascribed to increased concentration of Mo in soil solution as a result of its addition. Application of P also increased the amount of Mo in soil. Similar results were reported by Singh *et al.* (2014).

It could be concluded from the results that the application of 90 kg P₂O₅ ha⁻¹ and 3 kg Mo ha⁻¹ is beneficial in increasing the status of available N, P, K and Mo in soil besides crop yield and uptake of nutrients by the faba bean.

REFERENCES

- Jackson, M.L. (1973) Soil Chemical Analysis. Prentice Hall of India Private Limited, New Delhi.
- Singh, V., Ali, J. Singh, H. and Singh, J.P. (2016) Effect of sources of phosphorus and bio-fertilizers on yield nutrient uptake and quality of lentil. *Annals of Plant and Soil Research* **18**(1) : 14-17.
- Kushwaha, B. L. (1999) Studies on response of French bean to zinc, boron and molybdenum application. *Indian Journal of Pulses Research* **12** (1): 44-48
- Nadia, G. (2012) Influence of molybdenum on groundnut production under different nitrogen levels. *World Journal of Chemistry* **7** (2): 64-70.
- Srivastava, T. K. and Ahlawat I. P. S. (1995) Response of pea (*Pisum sativum* L.) to phosphorus, molybdenum and biofertilizers. *Indian Journal of Agronomy* **40**: 4, 630-635.
- Srivastava, T.K.; Ahlawat I.P.S. and Panwar J.D.S. (1998) Effect of phosphorus, molybdenum and bio-fertilizers on productivity of pea (*Pisum sativum* L.). *Indian Journal of Plant Physiology* **3** (3): 237.
- Alben Awomi T., Singh A. K., Manoj-Kumar and Bordoloi L. J. (2012) Effect of phosphorus, molybdenum and cobalt nutrition on yield and quality of mungbean (*Vigna radiata* L.) in acidic soil of northeast India. *Indian Journal of Hill Farming* **25**(2):22-26
- Nusakho Nyekha, Sharma, Y.K., Sharma, S.K. and Gupta, R.C. (2015) Effect of phosphorus and phosphorus solubilizing bacteria on performance of green gram and soil properties. *Annals of Plant and Soil Research* **17**(3): 323-325.
- Singh, S., Singh, H., Seema, Singh, J.P. and Sharma, V.K. (2014) Effect of integrated use of rock phosphate, molybdenum and phosphate solubilizing bacteria on lentil (*Lens culinaris*) in an alluvial soils. *Indian Journal of Agronomy* **59**(3): 433-438.
- Kumawat, B.L., Choudhary, A., Kumawat, A. and Kumawat, A. (2009) Effect of vermicompost and molybdenum on yield of green gram and fertility status of soil. *Annals of Plant and Soil Research* **11**: 92-95.